arXiv:1303.4858v2 [hep-ph] 25 Mar 2013

Constraints on Unparticles from $B_s \to \mu^+\mu^-$

Jong-Phil Lee*

Division of Quantum Phases & Devices, School of Physics,

Konkuk University, Seoul 143-701, Korea

Abstract

Unparticle contributions to the recently measured decay mode $B_s \to \mu^+\mu^-$ are analyzed. We consider only the scalar unparticles because vector unparticles are expected to provide negligible contributions. Assuming that the relevant coupling constants are real, we present allowed regions of coupling constants and the scaling dimension of the scalar unparticle. While the measured value of the branching ratio is very close to the standard model predictions, one cannot exclude the possible contributions from unparticles.

PACS numbers: 12.90.+b, 13.20.-v, 13.20.He

---

* jongphil7@gmail.com

Recently the LHCb collaboration reported the first evidence for the decay $B_s \to \mu^+\mu^-$ and an upper limit on $B_d \to \mu^+\mu^-$ as [1]

$$\mathrm{Br}(B_s \to \mu^+\mu^-) = \left(3.2^{+1.5}_{-1.2}\right) \times 10^{-9} \ , \tag{1}$$

$$\mathrm{Br}(B_d \to \mu^+\mu^-) < 9.4 \times 10^{-10} \ . \tag{2}$$

The result is quite consistent with the standard model (SM) predictions[2]

$$\mathrm{Br}(B_s \to \mu^+\mu^-)_{\mathrm{SM}} = (3.23 \pm 0.27) \times 10^{-9} \ , \tag{3}$$

$$\mathrm{Br}(B_d \to \mu^+\mu^-)_{\mathrm{SM}} = (1.07 \pm 0.10) \times 10^{-10} \ . \tag{4}$$

The decay of $B_s \to \mu^+\mu^-$ is very sensitive to new physics because in the SM the process occurs only through the loop contributions. However, it would be too early to declare that there are no new physics at all. Implications of the new observation in view of new physics can be found in [3–5]. In this paper, we examine the unparticle effects on $B_s \to \mu^+\mu^-$ decay.

Unparticle is a hypothetical concept associated with the scale invariance at high energy scales [6]. According to the unparticle scenario there is a scale-invariant hidden sector, and it couples to the SM particles very weakly at high energy scale $\Lambda_{\mathcal{U}}$. When seen at low energy, the hidden sector behaves in different ways from ordinary particles. That's the reason why the stuff is called as *unparticles*. In other words, unparticles behave like a fractional number of particles.

Consider a ultraviolet (UV) theory in the hidden sector at some high energy $\sim M_{\mathcal{U}}$ with the infrared (IR)-stable fixed point. It is quite convenient to describe the interaction between the UV theory and the SM sector in an effective theory formalism. Below the scale of $M_{\mathcal{U}}$, a UV operator $\mathcal{O}_{\mathrm{UV}}$ interacts with an SM operator $\mathcal{O}_{\mathrm{SM}}$ through $\mathcal{O}_{\mathrm{SM}}\mathcal{O}_{\mathrm{UV}}/M_{\mathcal{U}}^{d_{\mathrm{SM}}+d_{\mathrm{UV}}-4}$, where $d_{\mathrm{UV(SM)}}$ is the scaling dimension of $\mathcal{O}_{\mathrm{UV(SM)}}$. Through the renormalization flow, one can go down to a new scale $\Lambda_{\mathcal{U}}$. It appears through the dimensional transmutation where the scale invariance emerges. Below $\Lambda_{\mathcal{U}}$ the theory is matched onto the above interaction with the new unparticle operator $\mathcal{O}_{\mathcal{U}}$ as

$$C_{\mathcal{U}} \frac{\Lambda_{\mathcal{U}}^{d_{\mathrm{UV}}-d_{\mathcal{U}}}}{M_{\mathcal{U}}^{d_{SM}+d_{\mathrm{UV}}-4}} \mathcal{O}_{SM}\mathcal{O}_{\mathcal{U}} \ , \tag{5}$$

where $d_{\mathcal{U}}$ is the scaling dimension of $\mathcal{O}_{\mathcal{U}}$ and $C_{\mathcal{U}}$ is the matching coefficient. The value of $d_{\mathcal{U}}$ is not constrained to be integers because of the scale invariance. This unusual behavior of unparticles is reflected on the phase space of $\mathcal{O}_{\mathcal{U}}$. The spectral function of the unparticle is given by the two-point function of $\mathcal{O}_{\mathcal{U}}$ as

$$\begin{aligned}\rho_{\mathcal{U}}(P^2) &= \int d^4x\ e^{iP\cdot x}\langle 0|\mathcal{O}_{\mathcal{U}}(x)\mathcal{O}_{\mathcal{U}}^{\dagger}(0)|0\rangle \\ &= A_{d_{\mathcal{U}}}\theta(P^0)\theta(P^2)(P^2)^{d_{\mathcal{U}}-2}\ ,\end{aligned} \tag{6}$$

where

$$A_{d_{\mathcal{U}}} = \frac{16\pi^2\sqrt{\pi}}{(2\pi)^{2d_{\mathcal{U}}}}\frac{\Gamma(d_{\mathcal{U}}+\frac{1}{2})}{\Gamma(d_{\mathcal{U}}-1)\Gamma(2d_{\mathcal{U}})}\ , \tag{7}$$

is the normalization factor. The corresponding phase space is

$$d\Phi_{\mathcal{U}}(P) = \rho_{\mathcal{U}}(P^2)\frac{d^4P}{(2\pi)^4} = A_{d_{\mathcal{U}}}\theta(P^0)\theta(P^2)(P^2)^{d_{\mathcal{U}}-2}\frac{d^4P}{(2\pi)^4}\ , \tag{8}$$

and the propagator is given by

$$\int d^4x\ e^{iP\cdot x}\langle 0|T\mathcal{O}_{\mathcal{U}}(x)\mathcal{O}_{\mathcal{U}}^{\dagger}(0)|0\rangle = \frac{iA_{d_{\mathcal{U}}}}{2\sin d_{\mathcal{U}}\pi}\frac{e^{-i\phi_{d_{\mathcal{U}}}}}{(P^2+i\epsilon)^{2-d_{\mathcal{U}}}}\ , \tag{9}$$

where $\phi_{d_{\mathcal{U}}} = (d_{\mathcal{U}}-2)\pi$.

B physics is a good test bed for the unparticle effects [7, 8], including B_s-$\bar{B}_s$ mixing [9–12] (see also [13, 14] for meson mixing). One reason is that unparticles can contribute to the flavor changing neutral current at tree level.

For simplicity we only consider the left-handed currents coupled to scalar unparticles through

$$\mathcal{L}_{\mathcal{U}} = \frac{c_q}{\Lambda_{\mathcal{U}}^{d_{\mathcal{U}}}}\bar{q}'\gamma_\mu(1-\gamma_5)q\ \partial^\mu\mathcal{O}_{\mathcal{U}} + \frac{c_\ell}{\Lambda_{\mathcal{U}}^{d_{\mathcal{U}}}}\bar{\ell}'\gamma_\mu(1-\gamma_5)\ell\ \partial^\mu\mathcal{O}_{\mathcal{U}}\ , \tag{10}$$

where $c_{q,\ell}$ are dimensionless couplings. We assume that $c_{q,\ell}$ are real numbers. We do not consider vector unparticle contributions because they are expected to be highly suppressed. One can infer from Eq. (10) that the scalar unparticle contribution is proportional to $\sim (1/\Lambda_{\mathcal{U}}^2)^{d_{\mathcal{U}}}$, or more exactly (as will be shown later), $\sim (m_{B_s}^2/\Lambda_{\mathcal{U}}^2)^{d_{\mathcal{U}}}$. But the vector unparticle $\mathcal{O}_{\mathcal{U}}^\mu$ couples to the SM current as

$$\frac{c_V}{\Lambda_{\mathcal{U}}^{d_V-1}}\bar{q}'\gamma_\mu(1-\gamma_5)q\ \mathcal{O}_{\mathcal{U}}^\mu\ , \tag{11}$$

where d_V is the scaling dimension of $\mathcal{O}_{\mathcal{U}}^\mu$, and its contribution is $\sim (m_{B_s}^2/\Lambda_{\mathcal{U}}^2)^{d_V-1}$. But the unitarity constraints require that $d_{\mathcal{U}} \geq 1$ and $d_V \geq 3$ [15], resulting in much more suppression of the vector contribution [12].

$G_F = 1.16638 \times 10^{-5}$ GeV^{-2}	$\sin^2\theta_W = 0.23116$
$\alpha^{-1} = 127.937$	$V_{tb} = 0.999$
$\lvert V_{ts}\rvert = 0.0407$	$\phi_{ts} = -3.123$
$m_\mu = 105.658$ MeV	$m_t = 163.2$ GeV
$m_{B_s} = 5.3667$ GeV	$\tau_{B_s} = 1.497$ ps
$f_{B_s} = 234$ MeV	$\Lambda_{\mathcal{U}} = 1000$ GeV

TABLE I. Input parameters used in this paper. Here $\alpha^{-1} = \alpha(m_Z)^{-1}$, $m_t = m_t(m_t)$ in the $\overline{\rm MS}$ scheme, and ϕ_{ts} is the phase of V_{ts}.

The total decay rate of $B_s \to \mu^+\mu^-$ is now given by

$$\Gamma = \frac{1}{16\pi M_{B_s}}\sqrt{1-\frac{4m_\mu^2}{m_{B_s}^2}}\,\left|\langle\mu\mu|(\mathcal{H}_{eff}^{\rm SM}+\mathcal{H}_{eff}^{\mathcal{U}})|B_s\rangle\right|^2 , \tag{12}$$

where $\mathcal{H}_{eff}^{\rm SM}$ is the SM effective Hamiltonian, while the unparticle effective Hamiltonian $\mathcal{H}_{eff}^{\mathcal{U}}$ is

$$\mathcal{H}_{eff}^{\mathcal{U}} = \frac{A_{d_\mathcal{U}}e^{-i\phi_\mathcal{U}}}{\sin d_\mathcal{U}\pi}\left(\frac{m_{B_s}}{\Lambda_\mathcal{U}}\right)^{2d_\mathcal{U}}\left(\frac{m_\mu m_b}{m_{B_s}^4}\right)(c_q\cdot c_\ell)\left[\bar{b}(1-\gamma_5)s\right]\left[\bar{\ell}\gamma_5\ell\right] . \tag{13}$$

Now the branching ratio can be written as

$$\mathrm{Br}(B_s\to\mu\mu) = \mathrm{Br}_{\rm SM}\left(1+\frac{\mathcal{A}_\mathcal{U}}{\mathcal{A}_{\rm SM}}\right) . \tag{14}$$

Here $\mathrm{Br}_{\rm SM}$ is the SM prediction and

$$\mathcal{A}_\mathcal{U} = \frac{A_{d_\mathcal{U}}}{\sin d_\mathcal{U}\pi}\left(\frac{m_{B_s}}{\Lambda_\mathcal{U}}\right)^{2d_\mathcal{U}}(c_q\cdot c_\ell)\cos(\phi_\mathcal{U}+\phi_{ts}) \tag{15}$$

$$\mathcal{A}_{\rm SM} = \frac{G_F}{\sqrt{2}}\frac{\alpha}{2\sin^2\theta_W}m_{B_s}^2 Y(x_t)|V_{tb}^*V_{ts}| , \tag{16}$$

where $x_t = m_t^2/m_W^2$, $Y(x) = \eta_Y Y_0(x)$, and

$$Y_0(x) = \frac{x}{8}\left[\frac{x-4}{x-1}+\frac{3x}{(x-1)^2}\ln x\right] , \quad \eta_Y = 1.0113 . \tag{17}$$

In Table I, we summarize the input values used in this analysis. With the values of Table I, one gets $\mathrm{Br}_{\rm SM} = 3.54\times10^{-9}$, which is consistent with other literatures. To compare the theoretical prediction with the experimental result, one should consider the non-zero decay width effect of B_s meson [16–18]. According to [17],

$$\mathrm{Br}(B_s\to\mu^+\mu^-)_{\rm theo} = \left[\frac{1-y_s^2}{1+y_s\mathcal{A}_{\Delta\Gamma}}\right]\mathrm{Br}(B_s\to\mu^+\mu^-)_{\rm exp} , \tag{18}$$



FIG. 1. Allowed region in $d_{\mathcal{U}}$-c_f plane.

where

$$y_s \equiv \tau_{B_s}\frac{\Delta\Gamma_s}{2} = 0.088 \pm 0.014 \ . \tag{19}$$

Here

$$\mathcal{A}_{\Delta\Gamma} \equiv \frac{R_H - R_L}{R_H + R_L} \ , \tag{20}$$

where $R_{H(L)}\exp\left[-\Gamma^{(s)}_{H(L)}t\right]$ is the decay rate of the heavy (light) mass eigenstate. In our case of Eq. (13) (pseudoscalar leptonic operator) with real couplings, one can easily find that $\mathcal{A}_{\Delta\Gamma} = 1$ [18]. In light of Eq. (18), we should compare the theoretical prediction with

$$(1 - y_s)\mathrm{Br}(B_s \to \mu^+\mu^-)_{\mathrm{exp}} = (2.9^{+1.4}_{-1.1}) \times 10^{-9} \ . \tag{21}$$

Equation (21) is the reference value for our analysis.

As can be seen in Eqs. (13) and (15), the fermionic couplings of the scalar unparticle appear only in combination of $c_q \cdot c_\ell \equiv c_f$. Figure 1 shows the allowed values of c_f and $d_{\mathcal{U}}$ constrained by the measured branching ratio, Eq. (1). The behavior of Fig. 1 can be inferred from the Eqs. (14) and (15). Note that $\mathcal{A}_{\mathcal{U}}$ is proportional to $(m_{B_s}/\Lambda_{\mathcal{U}})^{2d_{\mathcal{U}}} \simeq (2.88\times10^{-5})^{d_{\mathcal{U}}}$, which suppresses the unparticle contribution to the total branching ratio significantly for $1 < d_{\mathcal{U}} < 2$. Thus for larger values of $d_{\mathcal{U}}$, the value of c_f can be large to fit the experimental result. As shown in Fig. 1, for $d_{\mathcal{U}} \gtrsim 1.4$, all the region of $0 \le c_f \le 1$ is allowed. That



FIG. 2. Allowed values of c_q and c_ℓ for $1.0 \le d_{\mathcal{U}} < 1.1$ (pink), $1.1 \le d_{\mathcal{U}} < 1.2$ (blue), $1.2 \le d_{\mathcal{U}} < 1.3$ (green), and $1.3 \le d_{\mathcal{U}} < 1.4$ (red). For $d_{\mathcal{U}} \gtrsim 1.4$, almost all the values of $0 \le c_{q,\ell} \le 1.0$ are allowed.

is the reason why we show only the case when $1 \le d_{\mathcal{U}} \le 1.4$ in Fig. 2. Figure 2 shows the allowed region of (c_q, c_ℓ) for different values of $d_{\mathcal{U}} \le 1.4$. The region is symmetric with respect to $c_q = c_\ell$ since the couplings only appear in form of $c_q \cdot c_\ell$. Note that the red points corresponding to $1.3 \le d_{\mathcal{U}} \le 1.4$ cover almost all the space of $0 \le c_{q,\ell} \le 1.0$. Current analysis is done for $\Lambda_{\mathcal{U}} = 1$ TeV. For larger values of $\Lambda_{\mathcal{U}}$, unparticle contribution gets smaller by $(m_{B_s}/\Lambda_{\mathcal{U}})^{2d_{\mathcal{U}}}$ and the allowed parameter space would become larger.

In conclusion, we have investigated the unparticle effects on $B_s \to \mu^+\mu^-$ decay. The experimental result is quite consistent with the SM, but it does not mean that there is no room for new physics. In this analysis only the scalar unparticles are considered because vector unparticles are expected to give negligible contributions. Assuming that scalar unparticles couple to the left-handed current, we provided the allowed regions of the couplings $c_{q,\ell}$ and the scaling dimension $d_{\mathcal{U}}$ for a fixed $\Lambda_{\mathcal{U}} = 1$ TeV. Since the unparticle contributions are proportional to $(m_{B_s}/\Lambda_{\mathcal{U}})^{2d_{\mathcal{U}}}$, the allowed parameter space of $c_{q,\ell}$ gets larger for large $d_{\mathcal{U}}$. The upper bound on $\mathrm{Br}(B_d \to \mu^+\mu^-)$ of Eq. (2) would not give strong constraints on the model parameters, since the SM prediction of Eq. (4) is almost an order of magnitude smaller. But if the branching ratio $\mathrm{Br}(B_d \to \mu^+\mu^-)$ is measured in near future, a combined analysis with

$B_s \to \mu^+\mu^-$ would give some hints on the flavor structure of unparticle interactions.

ACKNOWLEDGMENTS

The author thanks Kang Young Lee for his reminding of the subject and helpful discussions. This work is supported by WCU program through the KOSEF funded by the MEST (R31-2008-000-10057-0).

[1] RAaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **110**, 021801 (2013).

[2] A. J. Buras, J. Girrbach, D. Guadagnoli and G. Isidori, Eur. Phys. J. C **72**, 2172 (2012).

[3] A. Arbey, M. Battaglia, F. Mahmoudi and D. M. Santos, Phys. Rev. D **87**, 035026 (2013).

[4] D. Guadagnoli and G. Isidori, arXiv:1302.3909 [hep-ph].

[5] A. J. Buras, R. Fleischer, J. Girrbach and R. Knegjens, arXiv:1303.3820 [hep-ph].

[6] H. Georgi, Phys. Rev. Lett. **98**, 221601 (2007); Phys. Lett. B **650**, 275 (2007).

[7] C. H. Chen and C. Q. Geng, Phys. Rev. D **76**, 115003 (2007).

[8] R. Mohanta and A. K. Giri, Phys. Lett. B **660**, 376 (2008).

[9] R. Mohanta and A. K. Giri, Phys. Rev. D **76**, 075015 (2007).

[10] A. Lenz, Phys. Rev. D **76**, 065006 (2007).

[11] J. K. Parry, Phys. Rev. D **78**, 114023 (2008).

[12] J. -P. Lee, Phys. Rev. D **82**, 096009 (2010).

[13] X. Q. Li and Z. T. Wei, Phys. Lett. B **651**, 380 (2007).

[14] S. L. Chen, X. G. He, X. Q. Li, H. C. Tsai and Z. T. Wei, Eur. Phys. J. C **59**, 899 (2009).

[15] B. Grinstein, K. A. Intriligator and I. Z. Rothstein, Phys. Lett. B **662**, 367 (2008).

[16] S. Descotes-Genon, J. Matias and J. Virto, Phys. Rev. D **85**, 034010 (2012).

[17] K. De Bruyn, R. Fleischer, R. Knegjens, P. Koppenburg, M. Merk and N. Tuning, Phys. Rev. D **86**, 014027 (2012).

[18] K. De Bruyn, R. Fleischer, R. Knegjens, P. Koppenburg, M. Merk, A. Pellegrino and N. Tuning, Phys. Rev. Lett. **109**, 041801 (2012).